Item 5Cib BCWMC 10-17-13 Blue Water Science Report Appendices

# APPENDIX

- Appendix A: Notification of Fish Survey conducted by Blue Water Science
- Appendix B: Blue Water Science Sweeney Lake Fish Survey, 2013
- Appendix C: Blue Water Science Twin Lake Fish Survey, 2013
- Appendix D: MnDNR Sweeney Electrofishing Assessment, August 28, 2013
- Appendix E: MnDNR Sweeney and Twin Lakes Fish Surveys from 1991

## **APPENDIX A: Notification of Fish Survey**

From: Steve McComas [mailto:mccomas@pdink.com] Sent: Friday, September 13, 2013 8:26 AM To: Daryl Ellison ; Greg Salo Cc: Laura Jester Subject: Fish survey notification for Sweeney and Twin Lakes, Hennepin County

Hello all,

Blue Water Science will be conducting a fish survey in Sweeney Lake (MN ID 27-3501) and Twin Lake (MN ID 27-3502), Hennepin County, starting on Monday, September 16. We will set 8 fyke nets on Monday. The nets will be monitored daily on Tuesday and Wednesday and all fish will be weighed and measured and returned to the lake. The nets will be removed from the lake on Wednesday, September 18. The fish survey is sponsored by the Bassett Creek Watershed Management Commission with the objectives to determine if fish have an impact on water quality, to check for changes in the fish community structure since the last survey, and to supplement data collected with recent electrofishing that was conducted by the MnDNR – Fisheries..

This survey is being conducted under the permit number: 19414.

Thank you, Steve McComas BLUE WATER SCIENCE 550 South Snelling Avenue St. Paul, MN 55116 651 690 9602 mccomas@pclink.com



Trapnet Fish Survey for Sweeney Lake (27-0035-01), Hennepin County, September 17 - 19, 2013

Submitted to: Bassett Creek Watershed Management Commission



October 2013

Prepared by: Steve McComas Jo Stuckert Blue Water Science St. Paul, MN 55116

Appendix B - i

# Trapnet Fish Survey for Sweeney Lake, Hennepin County, September 17 - 19, 2013

## Introduction

Sweeney Lake (MnDNR ID: 27-0035-01) is a 66-acre lake and is connected by way of a channel to Twin Lake (MnDNR ID: 27-0035-02) which is 19 acres in area. In September of 2013, the Bassett Creek Watershed Management Commission contracted for trapnet fish surveys on both lakes with Blue Water Science. The fish surveys were conducted under MnDNR permit number 19414. The last trapnet surveys were conducted by the MnDNR in 1991. A recent electrofishing assessment was conducted on August 28, 2013.

The objectives of the trapnet survey were to characterize the fish community in both lakes and evaluate potential fish effects on lake water quality.

## Methods

Five standard trapnets were set in Sweeney Lake on September 17, 2013 and then were sampled daily on September 18 and 19, 2013. Each trapnet was a MnDNR style with two 4x6 foot square frames followed by two funnel mouth openings. A 50-foot lead net was staked on shore which led to the opening in the square frames. Net mesh size was 3/8 inch. Trapnet locations were shown in Figure 1. In Sweeney Lake, Net 5-1 was moved after the first day to a new location for the second day of sampling and the location is shown as Net 5-2 (Figure 1).



Figure 1. Locations of the trapnet sets on Sweeney Lake on September 17-19, 2013.

### Table 1. Summary of the Sweeney Lake fish survey conducted on September 18 and 19, 2013.

Species	SWEENEY Number of Fish per Net	MnDNR Normal Range
Black bullhead	1.1	1.3 - 26.0
Yellow bullhead	7.9	0.8 - 5.0
Black crappie	15	1.8 - 18.1
Bluegill	45	6.5 - 59.6
Carp	0.3	0.3 - 2.6
Gizzard shad	0	NA
Green sunfish	0	0.3 - 2.0
Hybrid sunfish	0.5	NA
Largemouth bass	0.2	0.3 - 0.8
Northern pike	0.8	NA
Pumpkinseed	1.6	0.8 - 5.3
White sucker	3.6	0.3 - 1.6
Yellow perch	0.3	0.3 - 1.5
Number of fish species	11	
Painted turtle	2.7	NA
Snapping turtle	0.2	NA
Softshell turtle	0.9	NA

#### Table 2. Sweeney Lake trapnet results for individual nets for the fish survey conducted in September, 2013.

Net	Black	Yellow	Black	Bluegill	Carp	Gizzard	Green	Hybrid	Large-		Pumpkin	White	Yellow		Turtles	
	bullhead	bullhead	crappie			shad	sunfish	sunfish	mouth bass	pike	seed	sucker	perch	Painted	Snap	Softshell
Tuesday (	(9/17/13)		1				1	I.	1	1				0		
1	3	4	21	48	1					3	3	10	1	1		1
2	1	12	43	134	1			3	1		4	2		3		
3	2	11	5	43	1			2		2	1	2				1
4	3	24	20	53							3	6	1	3		1
5			6	1								2				5
subtotal	9	51	95	279	3	0	0	5	1	5	11	22	2	7	0	8
average/ net	1.8	10	19	56	0.6	0	0	1.0	0.2	1.0	2.2	4.4	0.4	1.4	0	1.6
Wednesd	lay (9/18/13	3)														
1		5	20	56		4*				3	4	8		1		
2		3	15	46							1	2	1	3	1	
3		2	11	31					1			2		3	1	1
4	2	19	11	35								2				
5														13		
subtotal	2	29	57	168	0	4	0	0	1	3	5	14	1	20	2	1
average/ net	0.4	5.8	11	34	0	0.8	0	0	0.2	0.6	1.0	2.8	0.2	4.0	0.4	0.2
Total Fish (all nets)	11	80	152	447	3	4	0	5	2	8	16	36	3	27	2	9
Fish/ Trapnet	1.1	8.0	15	45	0.3	0.4	0	0.5	0.2	0.8	1.6	3.6	0.3	2.7	0.2	0.9

\*gizzard shad were regurgitated by Northern pike in the sample net.

### Table 3. Length frequency of fish species (as total length) for Sweeney and Twin Lakes fish survey for September 2013.

Length	Blac		Yello bullhe	w	Blac	k	Blueg	jill	Carı	D	Gree		Hybr		Largem		North		Pump		Whi		Yello	
(inches)	bullhe Sweeney	Twin	Sweeney	eaα Twin	crapp Sweeney		Sweeney	Twin	Sweeney	Twin	sunfi Sweeney		sunfi Sweeney		base Sweeney		pike Sweeney	e Twin	Sweeney		suck Sweeney	Twin	Sweeney	
<3							3	4			-								Í					
3					1		8																	
3.5					2		3								1				2					
4							3	1								1								<u> </u>
4.5 5							35 86	1 3				1	1						4	1			2	1
5 5.5							80 77	3 5					1						4				2	-
6					3		97	4					•	2					1	6				
6.5			1		18		74	2						1					1					
7			4		36	3	59	3					2										1	
7.5				1	48		4	3																<u> </u>
8			7		32	1	1						1	1										<u> </u>
8.5 9			4 10	2	5 3								1											+
9.5			10	1	2						-													-
10	3		10	1	1																			1
10.5	1		10																					
11	5		10	2	1																			$\perp$
11.5			7	1																				—
12	4		5	1																				┿
12.5 13	1 1		1	1																	1	-		<u> </u>
13.5	r		1	1																	1		1	+-
14				1																	· ·		1	+
14.5																					1			
15															1						6			
15.5																					6			<u> </u>
16																					8			<u> </u>
16.5 17																					5 7	1		+
17.5																					1			<u> </u>
18																								
18.5																								
19									1															
19.5																								<u> </u>
20																								<u> </u>
20.5 21																	1							+
21.5																								<u> </u>
22																								-
22.5																								
23									1															
23.5																								<u> </u>
24																						<u> </u>		—
24.5 25																							1	+
25.5																							1	+-
26									1								2				l		l	<u> </u>
26.5																								
27		I																						$\vdash$
27.5																								─
28 28.5																	1					<u> </u>		─
28.5 29																						-		+
29																					1	-		<u> </u>
30																	1						1	+
30.5																	1							
31																								<u> </u>
31.5					-						-							<u> </u>				<u> </u>		<u> </u>
32																		1				<u> </u>		<u> </u>
32.5 33																	1							+
33 33.5																	1							+-
33.5						-												1		1		-	1	<u> </u>
Total	11	0	79	12	152	4	447	22	3	0	0	1	5	4	2	1	8	1	16	7	36	1	3	1



## Trapnet Fish Survey for Twin Lake (27-0035-02), Hennepin County, September 17 - 19, 2013

Submitted to: Bassett Creek Watershed Management Commission



October 2013

Prepared by: Steve McComas Jo Stuckert Blue Water Science St. Paul, MN 55116

Appendix C - i

# Trapnet Fish Survey for Twin Lake, Hennepin County, September 17 - 19, 2013

## Introduction

Sweeney Lake (MnDNR ID: 27-0035-01) is a 66-acre lake and is connected by way of a channel to Twin Lake (MnDNR ID: 27-0035-02) which is 19 acres in area. In September of 2013, the Bassett Creek Watershed Management Commission contracted for trapnet fish surveys on both lakes with Blue Water Science. The fish surveys were conducted under MnDNR permit number 19414. The last trapnet surveys were conducted by the MnDNR in 1991. A recent electrofishing assessment was conducted on August 28, 2013.

The objectives of the trapnet survey were to characterize the fish community in both lakes and evaluate potential fish effects on lake water quality.

## Methods

Three standard trapnets were set in Twin Lake on September 17, 2013 and then were sampled daily on September 18 and 19, 2013. Each trapnet was a MnDNR style with two 4x6 foot square frames followed by two funnel mouth openings. A 50-foot lead net was staked on shore which led to the opening in the square frames. Net mesh size was 3/8 inch. Trapnet locations were shown in Figure 1. In Twin Lake, Nets 2-1 and 3-1 were moved after sampling on the first day to new locations and were sampled there on Day 2. The locations on Day 2 are shown as Net 2-2 and Net 3-2 (Figure 1).



Figure 1. Locations of the trapnet sets on Twin Lakes on September 17-19, 2013.

### Table 1. Summary of the Twin Lake fish survey conducted on September 18 and 19, 2013.

Species	TWIN Number of Fish per Net	MnDNR Normal Range
Black bullhead	0	1.3 - 26.0
Yellow bullhead	2.0	0.8 - 5.0
Black crappie	0.7	1.8 - 18.1
Bluegill	3.7	6.5 - 59.6
Carp	0	0.3 - 2.6
Gizzard shad	0	NA
Green sunfish	0.2	0.3 - 2.0
Hybrid sunfish	0.7	NA
Largemouth bass	0.2	0.3 - 0.8
Northern pike	0.2	NA
Pumpkinseed	0.7	0.8 - 5.3
White sucker	0.2	0.3 - 1.6
Yellow perch	0.2	0.3 - 1.5
Number of fish species	10	
Painted turtle	2.7	NA
Snapping turtle	0.3	NA
Softshell turtle	0	NA

### Table 2. Twin Lake trapnet results for individual nets for the fish survey conducted in September, 2013.

Net	Black	Yellow	Black	Bluegill	Carp	Gizzard	Green	Hybrid	Large-	North.	Pumpkin	White	Yellow		Turtles	
	bullhead	bullhead	crappie			shad	sunfish	sunfish	mouth bass	pike	seed	sucker	perch	Painted	Snap	Softshell
Tuesday (	(9/17/13)															
1				6			1				2				1	
2			4													
3		9	2	2									1			
subtotal	0	9	6	8	0	0	1	0	0	0	2	0	1	0	1	0
average/ net	0	3.0	2.0	27	0	0	0.3	0	0	0	0.7	0	0.3	0	0.3	0
Wednesd	lay (9/18/13	6)		r			r			r.			1			
1		6									3			1		
2																
3		3	1	12				4	1	1	2	1		15	1	
subtotal	0	9	1	12	0	0	0	4	1	1	5	1	0	16	1	0
average/ net	0	3.0	0.3	4.0	0	0	0	1.3	0.3	0.3	1.7	0.3	0	5.3	0.3	0
Total Fish (all nets)	0	18	7	20	0	0	1	4	1	1	7	1	1	16	2	0
Fish/ Trapnet	0	3.0	1.2	3.3	0	0	0.2	0.7	0.2	0.2	1.2	0.2	0.2	2.7	0.3	0

### Table 3. Length frequency of fish species (as total length) for Sweeney and Twin Lakes fish survey for September 2013.

Length	Blac		Yello bullhe	w	Blac	k	Blueg	jill	Carı	D	Gree		Hybr		Largem		North		Pump		Whi		Yello	
(inches)	bullhe Sweeney	Twin	Sweeney	eaα Twin	crapp Sweeney		Sweeney	Twin	Sweeney	Twin	sunfi Sweeney		sunfi Sweeney		base Sweeney		pike Sweeney	e Twin	Sweeney		suck Sweeney	Twin	Sweeney	
<3							3	4			-								Í					
3					1		8																	
3.5					2		3								1				2					
4							3	1								1								<u> </u>
4.5 5							35 86	1 3				1	1						4	1			2	1
5 5.5							80 77	3 5					1						4				2	-
6					3		97	4					•	2					1	6				
6.5			1		18		74	2						1					1					
7			4		36	3	59	3					2										1	
7.5				1	48		4	3																<u> </u>
8			7		32	1	1						1	1										<u> </u>
8.5 9			4 10	2	5 3								1											+
9.5			10	1	2						-													-
10	3		10	1	1																			1
10.5	1		10																					
11	5		10	2	1																			$\perp$
11.5			7	1																				—
12	4		5	1																				┿
12.5 13	1 1		1	1																	1	-		<u> </u>
13.5	r		1	1																	1		1	+-
14				1																	· ·		1	+
14.5																					1			
15															1						6			
15.5																					6			<u> </u>
16																					8			<u> </u>
16.5 17																					5 7	1		+
17.5																					1			<u> </u>
18																								
18.5																								
19									1															
19.5																								<u> </u>
20																								<u> </u>
20.5 21																	1							+
21.5																								<u> </u>
22																								-
22.5																								
23									1															
23.5																								<u> </u>
24																						<u> </u>		—
24.5 25																						<u> </u>	1	+
25.5																							1	+-
26									1								2				1		l	<u> </u>
26.5																								
27		I																						$\vdash$
27.5																								─
28 28.5																	1					<u> </u>		─
28.5 29																						-		+
29																					1	-		<u> </u>
30																	1						1	+
30.5																	1							
31																								<u> </u>
31.5					-						-							<u> </u>				<u> </u>		<u> </u>
32																		1				<u> </u>		<u> </u>
32.5 33																	1							+
33 33.5																	1							+-
33.5						-												1		1			1	<u> </u>
Total	11	0	79	12	152	4	447	22	3	0	0	1	5	4	2	1	8	1	16	7	36	1	3	1

## APPENDIX D: MnDNR Sweeney Electrofishing Assessment, August 28, 2013



Minnesota Department of Natural Resources Fisheries Management



STANDARD LAKE SURVEY REPORT

DRAFT VERSION - PRELIMINARY DATA (AS OF 08/29/2013)

#### Lake Name: Sweeney

DOW Number: 27-0035-01

Survey Type: Special Assessment Survey ID Date: 08/28/2013

			CIAL ASSESSM Electrofishing Quality Measur		
Lake Identifica	tion				1.1
	Alternate Lake Name: Primary Lake Class ID:	N/A 30	D	NR Sounding Map Number: Alternate Lake Class ID:	
Lake Location	×				
	Primary County:	Hennepin		Nearest Town:	Glenwood Junction
Legal Descripti	ions				
PLS	Lake Center: Section Lake Center:	Township - 29N 2902418	Range - 24W	Section - 18	
A	II Legal Descriptions: Hennepin County:	Township - 29N	Range - 24W	Sections - 18, 19	
Area Office					
	Area Name: Region Name:	Metro VVest Central		ORG Code: Region Number:	
Lake Access (Information b	ased on Re-Survey dat	ed 07/01/1991)	1.	1.00	
Station ID	Ownership	Public Use	Туре	Location / Comments	
	(Data exclu	ides records where publ	ic use is not designated	d or is designated "No Public Use")	
Lake Character	ristics				
	(planimetered acres):		GI	S Shoreline Length (miles):	2.11
	ilS Lake Area (acres):	67.64		Maximum Fetch (miles):	1.00
DC	OW Lake Area (acres):			etch Orientation (degrees):	
	Littoral Area (acres):			USGS Quad Map Number:	
	Area in MN (acres):	67.64 28.0	0	ISGS Quad 24K GIS Index:	3632
Ŋ	Maximum Depth (feet): Mean Depth (feet):				
	weart Deput (reet).	Tanca.			

Standard Lake Survey Report revision: 04/05/2011. Data Date: 08/29/2013 at 3:04 pm .

#### DRAFT

#### STANDARD LAKE SURVEY REPORT SPECIAL ASSESSMENT DATED 08/28/2013 FOR DOW NUMBER 27-0035-01

#### Watershed Characteristics

Major Watershed	Minor Watershed	
Name: Mississippi River-TC Watershed Number: 20 Watershed size (acres): 644,320	Name: Bassett Cr Watershed Number: 95 Watershed size (acres): 13,581	

#### Surveys And Investigations

Initial Survey:	08/03/1960.
Re-Survey:	07/01/1991, 07/13/1981.
Population Assessment:	07/02/1986, 07/13/1976.

#### Dissolved Oxygen And Temperature Profile Of Lake Water

Station ID	Sampling Date	Bottom Depth (Feet)	Sample Depth (Feet)	Water Temperature (°F)	Dissolved Oxygen (ppm)
VQ - 1	08/28/2013	22.0	Surface	82.6	7.2
			20	81.5	6.3
			4.0	80.6	6.1
			6.0	80.2	5.4
			8.0	80.1	4.9
			10.0	79.9	4.8
			12.0	79.7	4.2
			14.0	78.8	2.7
			16.0	78.1	1.4
			18.0	77.7	0.9
			20.0	76.8	0.5
			22.0	76.5	0.3

	Sampling	Sample	Secchi Depth	Field	Alkalinity		
Station ID	Date	Depth (Feet)	(Feet)	pH	(ppm)	Water Color	Color Cause
WQ = 1	08/28/2013	Surface	4.0	N\A	N/A	Brown Grn	Algae

2

#### Electrofishing Catch Summary for EF

Standard electrofishing

#### Total run-time for all stations: 01:37:00 Total on-time for all stations: 01:30:00 First Sampling Date: 08/28/2013 Last Sampling Date: 08/28/2013 Daylight Sampling: Yes Target Species: N/A

		Sum	mary By Num	bers	Summary By Weight (pounds)						
		Total	Number	per Hour	Total	Lbs pe	r Hour	Mean			
Abbr	Species	Number	Run-Time	On-Time	Weight	Run-Time	On-Time	Weight			
BLB	Black Bullhead	3	1.86	2.00	1.45	0.89	0.96	0.48			
BLC	Black Crappie	17	10.52	11.33	3.65	2.26	2.43	0.21			
BLG	Bluegill	86	53.20	57.33	9.97	6.17	6.65	0.12			
CAP	Common Carp	7	4.33	4.67	57.91	35.82	38.61	8.27			
GOS	Golden Shiner	5	3.09	3.33	0.48	0.30	0.32	0.10			
GSF	Green Sunfish	3	1.86	2.00	0.35	0.22	0.23	0.12			
HSF	Hybrid Sunfish	3	1.86	2.00	0.39	0.24	0.26	0.13			
LMB	Largemouth Bass	42	25.98	28.00	31.96	19.77	21.31	0.76			
NOP	Northern Pike	1	0.62	0.67	5.97	3,69	3.98	5.97			
PMK	Pumpkinseed	1	0.62	0.67	0.15	0.09	0.10	0.15			
WTS	White Sucker	8	4.95	5.33	13.05	8.07	8.70	1.63			
YEB	Yellow Bullhead	17	10.52	11.33	8.87	5.49	5.91	0.52			
YEP	Yellow Perch	2	1.24	1.33	0.08	0.05	0.05	0.04			

## DRAFT

## DRAFT

#### STANDARD LAKE SURVEY REPORT SPECIAL ASSESSMENT DATED 08/28/2013 FOR DOW NUMBER 27-0035-01

#### Length Frequency Distribution For EF

Standard electrofishing

(Field work conducted on 08/28/2013)

	BLB	BLC	BLG	CAP	GOS	GSF	HSF	LMB	NOP	PMK	WTS	YEB	YEP
< 3.00			1.1	-	1	~	-	2	-	1.1	-	1.1	-
3.00 - 3.49	÷.		1	- 51	1.0	-	-	1	1.1	10	-	÷.	
3.50 - 3.99			2		1	1	-	2		÷.	-	-	1.5
4.00 - 4.49	- n.		4	1.1	- C	1		1		1.1		-	1
4.50 - 4.99	-		24		-	1		2		1			1
5.00 - 5.49		19	23	1.1		1	1.151	-		1	- T	-	1.5
5.50 - 5.99		-	18	1.1	1	1	3	-		÷.			+
6.00 - 6.49			6		1			÷.					
6.50 - 6.99		1	7	÷.	1	÷	-	121	1.0	÷		÷	1.5
7.00 - 7.49		6	1			(÷	-	1		14	-	-	~
7.50 - 7.99	- ÷	9	E		1	+		÷.		+	-	- ÷	
8.00 - 8.49	1	1		1.1	~			2	1.11	(*)	-	1	1.1
8.50 - 8.99					-			1		÷	-	2	
9.00 - 9.49		-	1.00					2		÷.		3	1.10
9.50 - 9.99	1		-			e.,	-	3		- e	-	3	- ÷
10.00 - 10.49	1	1.0	-	-		-	10.0	4		-		2	1.0
10.50 - 10.99		-	1	-				4		. e	+	5	
11.00 - 11.49	1.1		-	1.1		-	1.4	4			1.4	1	1.12
11.50 - 11.99				-		~	-	2		÷	-		
12.00 - 12.99	1.4		.21	-			-	3	1.14				1.4
13.00 - 13.99			-	-		-	-	1		-	-		
14.00 - 14.99	1.1	- L.	- i		1			1		1.12	- 1		1
15.00 - 15.99		1.1	1	-			-	1			2		
16.00 - 16.99					- 1 i			1		1.2	6		
17.00 - 17.99	1.1		- L.	1.1	Q.			2	- D.	- C		1	1.0
18.00 - 18.99		1.4		1.2	- C.				1.1		1		1.2
19.00 - 19.99	2	1.2	- D		- 3.	12	1.0	1	- 0	S.		2	1.6.
20.00 - 20.99	- 2					62							- 1
21.00 - 21.99	2	1.2			10			1.1		- C.			1.1.5
22.00 - 22.99	- 2	- 2		1	- C			- 5		-		- 9	- 2
23.00 - 23.99	- Q.	- 5		2	1.0								
		- 2		-	- 3	- Č.		- 2		- 15			1.1
24.00 - 24.99				2									
25.00 - 25.99		19		1	- C	*		- 21			-	-	10
26.00 - 26.99				4		+		1					
27.00 - 27.99		1		4	~	+	-		- 24	+	-	-	
28.00 - 28.99				-		*			1.1		-		-
29.00 - 29.99	20	-			10		-		1		-		
30.00 - 30.99	۰.	1.2		7	1.0	7				- 5	-	₹.	1
31.00 - 31.99	<ul> <li></li> </ul>		-		× .	17						-	-
32.00 - 32.99	×.						-		-		-		
33.00 - 33.99	1	-	-	1.1	1.10	17	-	1.1	- B.		1.0	7	10
34.00 - 34.99				1.1	1.0	÷				÷	-	÷.	1.5
35.00 - 35.99	-	•			1.8		-	•		-		•	
= > 36.00	~		7		- 2	~	Ť	1	~	1	1	~	
	BLB	BLC	BLG	CAP	GOS	GSF	HSF	LMB	NOP	PMK	WTS	YEB	YEP
Total	3	17	88	7	5	3	3	40	1	1	8	17	2
Min. Length	8,15	6.57	3.27	22.24	2.68	4.84	5.71	2.56	29.33	5.31	15.12	8.46	4.13
Max. Length	10.04	8.43	15.16	27.36	7.80	5.63	5.87	17.32	29.33	5.31	16.93	11.02	4.72
The second s	9.30	7.53	5.48	24.93	5.87	5.26	5.77	9.86	29.33	5.31	16.24	9.90	4.43
Mean Length													
# Measured	3	17	88	7	5	3	3	40	1	1	8	17	2
No Lengths for	0	0	0	0	0	0	0	0	0	0	0	0	0

Note: Unless all fish were measured in the catch, totals shown for some length-frequency distributions may differ from the total number of fish in the catch, due to rounding of fractions used in the estimation of length frequency from a subsample of measured fish

4

DRA

5

#### Survey Crew Notes

Electrofishing assessment targeting all fish for Bassett Creek Watershed Management Commission

#### Discussion

The Bassett Creek Watershed Management Commission requested data on the fish community in Sweeney Lake and the connected Twin Lake. Specifically, the Commission was interested in the presence of common carp and gizzard shad. Since the most recent assessment was in 1991, an electrofishing assessment targeting all fish was conducted on Sweeney Lake during the day on August 28, 2013.

#### Sweeney Lake

Four transects, encompassing the entire lake, were electrofished. All transects were near shore in 4 feet of water or less. The conductivity of the lake water was high (1042 mS) and fish were shocked only moderately well. Despite this, in 1.5 h of electrofishing on-time, 195 fish were sampled, this included 13 different species. Bluegill, largemouth bass, black crappie, and yellow bullhead were the most abundant, respectively. Seven common carp were netted and measured. Many additional common carp were observed but were able to escape the electrical field before they were netted. One buffalo (Ictiobus sp.) was observed but could not be netted. Only netted fish are included in the survey report.

No gizzard shad were sampled in the 4 standard transects. However, areas of "rippling" water were observed off shore. Upon investigation with the electrofishing boat, these "ripples" were caused by schools of gizzard shad. The water was calm and these schools were observed in many areas throughout the lake. All gizzard shad that were shocked ranged from 3 to 5 inches. Since the shad were not sampled in the standard transects, they are not included in the survey report. Gizzard shad are not common in lakes of this type but they seemed relatively abundant in Sweeney Lake.

#### Twin Lake

The channel between Sweeney and Twin Lakes is shallow. At times the electrofishing boat had to be propelled by a push pole or crew members waded in the water and pushed the boat by hand. There was approximately 6 inches of clearance between the boat rails and the top of the bridge that leads to Twin Lake. During high water it may be difficult to pass under this bridge.

Once in Twin Lake, it was immediately obvious that the lake is heavily used by the public to recreate. Many people had accessed the lake at several points along the eastern shore and were swimming. Due to the number of people swimming laps in this 19-acre lake, it was determined that it was unsafe to use electricity to sample fish. No fish sampling was conducted. Water clarify was noticeably greater than Sweeney Lake and common carp and bluegill were visually observed.

DRAFT

STANDARD LAKE SURVEY REPORT SPECIAL ASSESSMENT DATED 08/28/2013 FOR D.OW NUMBER 27-0035-01

Approval Dates And Notices

Date Approved By Metro West Area Fisheries Supervisor:

Date Approved By Central Region Fisheries Manager:

This Draft version of the Standard Lake Survey Report contains preliminary data (as of 08/29/2013), and is therefore subject to change at any time.



Minnesota Department of Natural Resources

By accepting the data in this report, the user agrees the data will be used for personal benefit and not for profit. Any other uses or publication of the data needs the consent of the Department. The Minnesota Department of Natural Resources assumes no responsibility for actual or consequential damage incurred as a result of any user's reliance on the data.

Standard Lake Survey Report revision: 04/05/2011-RJE. Data Date: 08/29/2013 at 3:04 pm .

#### REPORT OVERVIEW - FOR OFFICE USE ONLY

(This page is not part of the Standard Lake Survey Report and should be discarded )

#### Lake Name: Sweeney

#### DOW Number: 27-0035-01

### Survey Type: Special Assessment

Survey ID Date: 08/28/2013

#### Electrofishing, Water Quality Measurement

#### Survey Status: Proposed

The following 22 (of 31) report components are not included in this report.

- 1. Current Water Level
- 2 Benchmark And Gauge Descriptions / Locations
- 3. Water Level History\*
- 4. Water Level History Readings\*
- 5. Water Level History Station Summary\*
- 6. Lake Inlets
- 7. Additional Inlet Information
- 8. Lake Outlets
- 9. Additional Outlet Information
- 10. Water Control Structure (Dam)
- 11. Surrounding Watershed Characteristics, Shoreline Characteristics, and Riparian Landscape Observations
- 12. Resorts And Campgrounds
- 13. Fish Spawning Conditions
- 14. Erosion And Pollution
- 15. Fish Diseases And Parasites
- 16. Aquatic Vegetation And Shoalwater Substrates
- 17. Laboratory Analysis Of Water Chemistry
- 18. Length At Capture With Last Incremental Length\*
- 19. Back-Calculated Lengths
- 20. Age Class Frequency Distributions
- 21. Other Species (added to revision 03/24/2009)
- 22. Water Quality (Winter Observations) (added to revision 01/21/2010)

\* Water Level History report: This data has not yet been migrated into the Fisheries LSM database. On 01/08/2009, two additional Water Level History report components (Readings and Station Summary) were added.

\* Length At Capture With Last Incremental Length report. The following criteria must be met for a report to be generated:

- 1. The fish species must have an assigned body scale constant.
- 2. Fish must have an "official" age assigned.
- 3. Fish must have a digitized measurement marked for back calculation use.

Note: The data source for Length and Age Class Frequency Distribution tables is updated twice daily - once at noon and once overnight. Any changes to the data made after noon on 08/29/2013 may not be reflected in the Distribution tables until 08/30/2013.

Fisheries Lake ID = 3129, Survey ID = 5171323833038000

Standard Lake Survey Report revision: 04/05/2011. Data Date: 08/29/2013 at 3:04 pm .

## APPENDIX E: MnDNR Sweeney and Twin Lakes Fish Surveys from 1991

Minnesota Department of Natural Resources

# Lake information report

## Name: SWEENEY

Nearest Town: GOLDEN VALLEY Primary County: Hennepin Survey Date: 07/01/1991 Inventory Number: 27-0035-01

## Lake Characteristics

Lake Area (acres): 66.00 Littoral Area (acres): 34.00 Maximum Depth (ft): 28.00 Water Clarity (ft): N/A Dominant Bottom Substrate: N/A Abundance of Aquatic Plants: N/A Maximum Depth of Plant Growth (ft): N/A

## Fish Sampled for the 1991 Survey Year

		Number of	fish per net	Average	Normal	
Species	<u>Gear Used</u>	Caught	Normal Range	Fish Weight (lbs)	Range (lbs)	
White Sucker	Gill net	1.0	0.5 - 2.0	1.40	1.0 - 2.2	
White Crappie	Gill net	1.0	0.5 - 4.8	0.65	0.2 - 0.3	
Northern Pike	Gill net	2.0	2.5 - 7.9	3.55	1.8 - 3.3	
Largemouth Bass	Gill net	1.0	0.3 - 1.1	1.10	0.4 - 1.5	
Common Carp	Gill net	2.5	0.5 - 4.0	4.58	1.0 - 3.2	
<u>Bluegill</u>	Gill net	21.5	N/A - N/A	0.12	N/A - N/A	
Black Crappie	Gill net	4.5	1.9 - 18.0	0.22	0.1 - 0.3	
Black Bullhead	Gill net	7.0	5.2 - 56.2	0.35	0.2 - 0.5	
Yellow Perch	Trap net	0.2	0.3 - 1.5	0.10	0.1 - 0.2	
Yellow Bullhead	Trap net	2.8	0.8 - 5.0	0.63	0.4 - 0.7	
White Sucker	Trap net	3.7	0.3 - 1.6	1.26	1.0 - 2.2	
Smallmouth Buffalo	Trap net	1.5	N/A - N/A	2.07	N/A - N/A	
Pumpkinseed Sunfish	Trap net	4.0	0.8 - 5.3	0.08	0.1 - 0.2	
Northern Pike	Trap net	1.5	N/A - N/A	3.08	N/A - N/A	
Largemouth Bass	Trap net	0.3	0.3 - 0.8	0.60	0.2 - 1.1	
Hybrid Sunfish	Trap net	0.5	N/A - N/A	0.10	N/A - N/A	
Green Sunfish	Trap net	0.7	0.3 - 2.0	0.10	0.1 - 0.1	
Common Carp	Trap net	4.8	0.3 - 2.6	4.12	2.0 - 4.5	
<u>Bluegill</u>	Trap net	124.5	6.5 - 59.6	0.08	0.1 - 0.2	
Black Crappie	Trap net	0.5	1.8 - 18.1	0.33	0.2 - 0.3	
Black Bullhead	Trap net	11.0	1.3 - 26.0	0.35	0.2 - 0.5	

Normal Ranges represent typical catches for lakes with similar physical and chemical characteristics.

## Length of Selected Species Sampled for All Gear for the 1991 Survey Year

Species	Number of fish caught in each category (inches)									
Species	0-5	6-8	9-11	12-14	15-19	20-24	25-29	30+	Total	
Yellow Perch	1	0	0	0	0	0	0	0	1	
Yellow Bullhead	0	5	7	5	0	0	0	0	17	
White Crappie	0	1	0	1	0	0	0	0	2	
Pumpkinseed Sunfish	24	0	0	0	0	0	0	0	24	
Northern Pike	0	0	0	0	3	4	5	1	13	
Largemouth Bass	1	0	1	2	0	0	0	0	4	
Hybrid Sunfish	3	0	0	0	0	0	0	0	3	
Green Sunfish	4	0	0	0	0	0	0	0	4	
<u>Bluegill</u>	233	38	0	0	0	0	0	0	271	
Black Crappie	0	12	0	0	0	0	0	0	12	
Black Bullhead	2	55	23	0	0	0	0	0	80	

## Status of the Fishery (as of 07/01/1991)

THE PRESENT FISH POPULATION IS DOMINATED BY BLHS, SMALL CEN, CATOSTOMIDS AND CYPRINIDS. BLGS AND BLBS WERE SAMPLED WELL ABOVE MEDIAN LEVELS; BLG AVERAGED 12.06 FISH PER POUND AND HAVE A PSD=10.1. BLBS AVERAGED 8.5-8.9 INCHES TOTAL LENGTH AND 2.88 FISH PER POUND. CARP WERE SAMPLED SLIGHTLY ABOVE MEDIAN LEVELS FOR NUMBER, BUT WELL ABOVE MEDIAN LEVELS FOR POUNDS PER SET FOR GILLNETS. CARP WERE SAMPLED WELL ABOVE TRAPNET MEDIANS. CAPTURED CARP AVERAGED 20.0 INCHES TOTAL LENGTH AND 4.19 POUNDS PER INDIVIDUAL. WTS AND NOP WERE SAMPLED NEAR MEDIAN LEVELS FOR GILLNETS AND WELL ABOVE MEDIAN LEVELS FOR TRAPNETS. SAMPLED NOP EXHIBITED A CALCULATED RSD = 76.92% FOR COMBINED GEAR TYPES. NOP GROWTH RATE APPEARS TO BE SLIGHTLY BELOW THE STATEWIDE AVERAGE. STOCKING OF LMB BY PRIVATE PERMIT ISSUED TO THE SWEENEY LAKE ASSOCIATION IS CONTRIBUTING ANOTHER PREDATOR SPECIES TO THE OVERALL POPULATION. GROWTH RATE FOR INDIVIDUALS SAMPLED IS SLIGHTLY ABOVE AVERAGE AND SEVERL YOY LMB WERE CAPTURED DURING SHORELINE SEINING. BLC WERE SAMPLED ABOVE MEDIAN LEVELS FOR GILLNETS, BUT BELOW FOR TRAPNETS. WHC AND YEP WERE ALSO CAPTURED DURING THIS SURVEY, BUT IN LOW NUMBERS. SEVERAL SAB WERE SAMPLED FOR THE FIRST TIME SINCE THE INITIAL SURVEY IN 1960. YEBS ARE ALSO PRESENT AND WERE SAMPLED ABOVE MEDIAN LEVELS FOR NUMBER OF INDIVIDUALS PER TRAPNET SET.



# Lake information report

## Name: SWEENEY TWIN

Nearest Town: GOLDEN VALLEY Primary County: Hennepin Survey Date: 07/02/1991 Inventory Number: 27-0035-02

### **Public Access Information**

Ownership	Туре	Description
Unknown	Unknown	ACCESS GAINED THROUGH GOLDEN VALLEY HEALTH CENTER PROPERTY BETWEEN SWEENEY

## Lake Characteristics

Lake Area (acres): 19.00	
Littoral Area (acres): 8.00	
Maximum Depth (ft): 56.00	
Water Clarity (ft): N/A	

Dominant Bottom Substrate: N/A Abundance of Aquatic Plants: N/A Maximum Depth of Plant Growth (ft): N/A

## Fish Sampled for the 1991 Survey Year

		Number of	fish per net	Average Fich	Normal	
Species	<u>Gear Used</u>	Caught	<u>Normal</u> <u>Range</u>	<u>Average Fish</u> <u>Weight</u> (Ibs)	<u>Range</u> (lbs)	
Yellow Bullhead	Gill net	0.5	1.0 - 6.9	0.20	0.4 - 0.7	
White Sucker	Gill net	0.5	0.5 - 2.0	0.30	1.0 - 2.2	
Pumpkinseed Sunfish	Gill net	1.5	N/A - N/A	0.07	N/A - N/A	
Northern Pike	Gill net	0.5	2.5 - 7.9	1.10	1.8 - 3.3	
<u>Bluegill</u>	Gill net	2.5	N/A - N/A	0.10	N/A - N/A	
Black Bullhead	Gill net	7.0	5.2 - 56.2	0.34	0.2 - 0.5	
Yellow Bullhead	Trap net	1.2	0.8 - 5.0	0.42	0.4 - 0.7	
White Sucker	Trap net	0.4	0.3 - 1.6	1.00	1.0 - 2.2	
Smallmouth Buffalo	Trap net	0.6	N/A - N/A	1.73	N/A - N/A	
Pumpkinseed Sunfish	Trap net	1.4	0.8 - 5.3	0.16	0.1 - 0.2	
Northern Pike	Trap net	0.2	N/A - N/A	2.50	N/A - N/A	
Largemouth Bass	Trap net	0.4	0.3 - 0.8	0.15	0.2 - 1.1	
Hybrid Sunfish	Trap net	1.4	N/A - N/A	0.20	N/A - N/A	
Green Sunfish	Trap net	0.2	0.3 - 2.0	0.10	0.1 - 0.1	
Golden Shiner	Trap net	0.2	0.2 - 1.4	0.35	0.1 - 0.1	
Common Carp	Trap net	0.8	0.3 - 2.6	3.30	2.0 - 4.5	
<u>Bluegill</u>	Trap net	29.8	6.5 - 59.6	0.13	0.1 - 0.2	
Black Crappie	Trap net	1.4	1.8 - 18.1	0.30	0.2 - 0.3	
Black Bullhead	Trap net	3.0	1.3 - 26.0	0.33	0.2 - 0.5	

Normal Ranges represent typical catches for lakes with similar physical and chemical characteristiCS.

## Length of Selected Species Sampled for All Gear for the 1991 Survey Year

Species	Number of fish caught in each category (inches)									
Species	0-5	6-8	9-11	12-14	15-19	20-24	25-29	30+	Total	
Yellow Bullhead	0	3	4	0	0	0	0	0	7	
Pumpkinseed Sunfish	9	1	0	0	0	0	0	0	10	
Northern Pike	0	0	0	0	1	1	0	0	2	
Largemouth Bass	2	0	0	0	0	0	0	0	2	
Hybrid Sunfish	6	1	0	0	0	0	0	0	7	
<u>Green Sunfish</u>	1	0	0	0	0	0	0	0	1	
<u>Bluegill</u>	68	15	0	0	0	0	0	0	83	
Black Crappie	0	7	0	0	0	0	0	0	7	
Black Bullhead	0	24	5	0	0	0	0	0	29	

## Status of the Fishery (as of 07/02/1991)

THE PRESENT FISH POPULATION IS DOMINATED BY BLHS AND SMALL CEN. ONLY BLGS WERE SAMPLED ABOVE ALL MEDIAN LEVELS FOR BOTH GEAR TYPES. CAPTURED BLGS AVERAGED 7.86 FISH PER POUND AND EXHIBITED A CALCULATED PSD=16.7%. CARP, BLC, AND YEB WERE SAMPLED BELOW MEDIAN LEVELS FOR TRAPNETS. PMK WERE SAMPLED BELOW MEDIAN LEVELS FOR TRAPNETS, BUT ABOVE THE MEDIANS FOR GILLNETS. BLB WERE SAMPLED IN THE SECOND HIGHEST ABUNDANCE AND ABOVE MEDIAN LEVELS FOR GILLNETS, BUT BELOW LOCAL MEDIAN LEVELS FOR TRAP NETS. OTHER SPECIES SAMPLED, BUT IN LOW ABUNDANCE, INCLUDE NOP, WTS, GOS, HSF, GSF, AND LMB. LMB SHOWED AVERAGE GROWTH FOR THE AGE 1+ INDIVIDUALS SAMPLED AND YOY BASS WERE SAMPLED DURING SHORELINE SEINING. SAB WERE CAPTURED FOR THE FIRST TIME SINCE FISH SURVEYS BEGAN IN 1960. THE ONLY FISH SPECIES SAMPLED IN SWEENEY'S TWIN THAT WAS NOT CAPTURED IN SWEENEY LAKE WAS GOS. WHC AND YEP WERE SAMPLED IN SWEENEY LAKE, BUT NOT IN SWEENEY'S TWIN.